

Boundness of b_2 for hyperkähler manifolds with vanishing odd-Betti numbers

Nikon Kurnosov¹

ABSTRACT. We prove that b_2 is bounded for hyperkähler manifolds with vanishing odd-Betti numbers. The explicit upper boundary is conjectured. Following the method described by Sawon we prove that b_2 is bounded in dimension eight and ten in the case of vanishing odd-Betti numbers by 24 and 25 respectively.

1 Introduction

A Riemannian manifold (M, g) is called hyperkähler if it admits a triple of a complex structures I, J, K satisfying quaternionic relations and Kähler with a respect to g . A hyperkähler manifold is always holomorphically symplectic. By the Yau Theorem [Y], a hyperkähler structure exists on a compact complex manifold if and only if it is Kähler and holomorphically symplectic.

Definition 1.1: A compact hyperkähler manifold M is called *simple* if $\pi_1(M) = 0$, $H^{2,0}(M) = \mathbb{C}$.

By Bogomolov decomposition theorem [B] any hyperkähler manifold admits a finite covering which is a product of a torus and several simple hyperkähler manifolds. Huybrechts [H] has proved finiteness of the number of deformation classes of holomorphic symplectic structures on each smooth manifold. In most dimensions we only know two examples of simple hyperkähler manifolds due to Beauville [Bea]. This two infinite series are the Hilbert schemes of $K3$ and the generalized Kummer varieties [Bea]. Except them two sporadic examples of O’Grady in dimension six and ten are known [O1, O2]. It is known [H] that there are only finitely many families with a given second cohomology lattice, hence the bounds on second Betti number provide restrictions on the number of deformation classes of hyperkähler manifolds. There is Beauville Conjecture [Bea1]

¹The article was prepared within the framework of a subsidy granted to the HSE by the Government of the Russian Federation for the implementation of the Global Competitiveness Program. Nikon Kurnosov is partially supported by grant MK-1297.2014.1., and RScF grant, project 14-21-00053 dated 11.08.14.

Conjecture 1.2: The number of deformation types of compact irreducible hyperkähler manifolds is finite in any dimension (at least for given b_2).

In the complex dimension four, Guan [G] proved that the second Betti number of a simple compact hyperkähler manifold of dimension four is bounded above by 23. Guan also have proved boundary conditions for b_3 using Rozansky-Witten invariants [HS]. In dimension six inequality from Rozansky-Witten invariants could also be obtained in dimension six [K].

In dimension six the boundary conditions for b_2 has been obtained by Sawon [S]. In this paper we generalize Sawon's result on Hodge diamond structure for dimensions eight and ten.

Acknowledgments. Author is grateful to his supervisor Misha Verbitsky for discussions, also author wishes to thank Justin Sawon who kindly describe his method, and Fedor Bogomolov for some ideas. Author would like thank Evgeny Shinder and Tom Bridgeland for the possibility to give a talk on this subject at Sheffield University.

2 Preliminaries

In this Section we recall main properties of cohomology groups of hyperkähler manifolds.

There is the following theorem of Verbitsky [V1]

Theorem 2.1: Let M be an irreducible hyperkähler manifold of complex dimension a $2n$ and let $SH^2(M, \mathbb{C}) \subset H^*(M, \mathbb{C})$ be the subalgebra generated by $H^2(M, \mathbb{C})$. Then $SH^2(M, \mathbb{C}) = S^*H^2(M, \mathbb{C}) / \langle \alpha^{n+1} \mid q(\alpha) = 0 \rangle$, where q is a Beaville-Bogomolov-Fujiki-form.

The inclusion $S^n H^2(M) \hookrightarrow H^{2n}$ follows from this Theorem.

It is known that in a hyperkähler case there is an action of $\mathfrak{so}(5)$ -algebra [V1], and moreover an action of $\mathfrak{so}(b_2 + 2, \mathbb{C})$ described by Looijenga and Lunts [LL], and Verbitsky [V2]. By this results one can decompose $H^{even}(M, \mathbb{C})$ into irreducible representations for this $\mathfrak{so}(b_2 + 2, \mathbb{C})$ -action. The Hodge diamond is the projection onto a plane of the (higher-dimensional)

weight lattice of $\mathfrak{so}(b_2 + 2, \mathbb{C})$ so we can figure out the highest weights which lie in octant of Hodge diamond.

Now irreducible representations are isomorphic to $\Lambda^i \mathbb{C}^{b_2+2}$. Thus, their dimensions are polynomial in terms of b_2 . Contribution of $\Lambda^i \mathbb{C}^{b_2+2}$ to each even cohomology groups also has dimension polynomial in terms of b_2 .

In early 90-s Salamon [Sa] has used hyperkähler symmetries of Hodge numbers and expression of Euler characteristic to prove the following equation with Betti numbers:

$$nb_{2n} = 2 \sum_{i=1}^{2n} (-1)^i (3i^2 - n) b_{2n-i}.$$

3 Boundary conditions for b_2

In this Section we prove the following

Theorem 3.1: Let M be a hyperkähler manifold of complex dimension $2n$ with vanishing odd-Betti numbers. Then the second Betti number is bounded.

Proof:

Consider the $2k$ -th Betti number. There is inclusion of $\mathrm{Sym}^k(H^2)$ into $H^{2k}(M)$. Thus, its contribution is polynomial of degree k . And also there are contributions of several $\mathfrak{so}(b_2 + 2, \mathbb{C})$ -modules into $H^{2k}(M)$. The dimension of each of them is polynomial in terms of b_2 , and the contribution of $\Lambda^i \mathbb{C}^{b_2+2}$ in $H^{2k}(M)$ is polynomial of degree at most $k - 1$. Recall that the highest weights lie in octant of Hodge diamond, and denote them as c, d and etc. For instance, c is dimension of $h_{prim}^{3,1}$.

Then for all even Betti numbers one could write that

$$b_{2k} = \frac{1}{(k-1)!} \prod_{i=0}^{i=k-1} (b_2 + i) + P_k(b_2, c, d, e, f, \dots),$$

where first term correspond to the dimension of symmetric power, and the second one – to all contributions of irreducible $\mathfrak{so}(b_2 + 2, \mathbb{C})$ -modules into $H^{2k}(M)$.

From Salamon's equation

$$nb_{2n} = 2 \sum_{i=1}^{2n} (-1)^i (3i^2 - n) b_{2n-i}.$$

using vanishing of odd-Betti numbers we obtain the following

$$\begin{aligned} & \frac{1}{(n-1)!} \prod_{i=0}^{i=n-1} (b_2 + i) + P_n(b_2, c, d, e, f, \dots) = \\ & = \sum_{j=1, j \text{ even}}^{j=2n} \left[(3j^2 - n) \frac{1}{(n-j/2)!} \prod_{i=0}^{i=n-j/2-1} (b_2 + i) + P_j(b_2, c, d, e, f, \dots) \right]. \end{aligned}$$

Rearranging terms in equation above (we put terms corresponding to contributions of symmetric powers of H^2 on the left-hand side and all others on the right-hand side):

$$\begin{aligned} & -\frac{1}{(n-1)!} \prod_{i=0}^{i=n-1} (b_2 + i) + \sum_{j=1, j \text{ even}}^{j=2n} (3j^2 - n) \frac{1}{(n-j/2)!} \prod_{i=0}^{i=n-j/2-1} (b_2 + i) = \\ & = P_n(b_2, c, d, e, f, \dots) - \sum_{j=1, j \text{ even}}^{j=2n} P_j(b_2, c, d, e, f, \dots). \end{aligned}$$

Denote by b_2^l the maximal root (in terms of b_2) of polynomial $P(b_2, n)$ on the left-hand side. Then the polynomial on the left-hand side is negative than $b_2 \geq b_2^l$. This polynomial obviously has some positive roots since it is positive for $b_2 = 0$.

The numbers c, d, e, f, \dots are positive since they are numbers of generators of irreducible $\mathfrak{so}(b_2 + 2, \mathbb{C})$ -modules. The leading coefficient of polynomial $Q_n(b_2, c, d, e, f, \dots)$ on the right-hand side is positive. Thus, this polynomial is positive then $b_2 \geq b_2^r$, where b_2^r is the maximal root of $Q_n(b_2, c, d, e, f, \dots)$.

Hence we get $b_2 \leq \max[b_2^l, b_2^r]$. In the case if right-hand side polynomial $Q_n(b_2, c, d, e, f, \dots)$ does not have any roots, then $b_2 \leq b_2^l$. ■

Conjecture 3.2: The second Betti number b_2 is bounded by the maximal root of the following polynomial

$$P(b_2, n) = -\frac{1}{(n-1)!} \prod_{i=0}^{i=n-1} (b_2 + i) + \sum_{j=1, j \text{ even}}^{j=2n} (3j^2 - n) \frac{1}{(n-j/2)!} \prod_{i=0}^{i=n-j/2-1} (b_2 + i),$$

which is denoted as b_2^l .

To prove this conjecture one need to check that $Q_n(b_2, c, d, e, f, \dots)$ is positive than $b_2 \geq b_2^l$. Author does not know the explicit proof of this Conjecture 3.2.

Proposition 3.3: In conditions of Theorem 3.1 we have

$$P(b_2, n) = -\frac{1}{n!} \left(\prod_{i=3}^{i=n-1} (b_2 + i) \right) \cdot (b_2 + 2n) \cdot (b_2^2 - 21b_2 + 2 - 96n),$$

and

$$b_2^l = \frac{21 + \sqrt{433 + 96n}}{2}.$$

Proof:

We can see that last four terms of $P(b_2, n)$ are divided by $(b_2 + 3)$ and factor is

$$\frac{1}{3}((12n^2 - 73n + 108)b_2^2 + 3(12n^2 - 49n + 48)b_2 + 12n(n - 1))$$

Then, we can claim by induction (we omit explicit calculations) that sum of last k terms of $P(b_2, n)$ is

$$2 \frac{1}{(k-1)!} \left(\prod_{i=3}^{i=k-1} (b_2 + i) \right) \cdot (A_k \cdot b_2^2 + 3B_k \cdot b_2 + 12n(n-1)),$$

where $A_k = (12n^2 - (73 + 24(k-4))n + 108 + 60k + 24 \frac{(k-4)(k-3)}{2})$, and $B_k = (12n^2 - (49 + 16k)n + 48 + 24k + 8 \frac{(k-4)(k-3)}{2})$.

Thus, we can write our polynomial $P(b_2, n)$ in the following form

$$P(b_2, n) = \frac{1}{(n-1)!} \prod_{i=0}^{i=n-1} (b_2 + i) + 2 \frac{1}{(n-1)!} \left(\prod_{i=3}^{i=n-1} (b_2 + i) \right) \cdot (A_n \cdot b_2^2 + 3B_n \cdot b_2 + 12n(n-1)).$$

After rearranging of common factors we get the statement. ■

4 Dimension eight and ten

To find explicit boundary condition we shall use the method described by Justin Sawon [S]. In his work he has studied how different $\mathfrak{so}(b_2 + 2, \mathbb{C})$ -modules sit inside Hodge diamond.

Theorem 4.1: Let M be a eight-dimensional hyperkähler manifold with $H^{2k+1}(M, \mathbb{C}) = 0$. Then $b_2 \leq 24$.

Proof:

We will write some explicit splitting of b_4, b_6 , and b_8 in terms of b_2 and generators of primitives part of cohomologies.

There is an action of $\mathfrak{so}(b_2 + 2, \mathbb{C})$ on the complex cohomology of M . Under this action, an element of $H_{\text{pr}}^{3,1}(M)$ will generate an irreducible $\mathfrak{so}(b_2 + 2, \mathbb{C})$ -module of dimension $\frac{(b_2+2)(b_2+1)b_2}{6}$. In Hodge diamond this part sits as

[illegible]

In the same way we have an irreducible $\mathfrak{so}(b_2 + 2, \mathbb{C})$ -module generated by additional elements of $H_{\text{pr}}^{2,2}$, which sits inside the Hodge diamond as

[illegible]

Recall that $b_4 = \frac{(b_2+1)b_2}{2} + cb_2 + d$, where d – part of $H_{\text{pr}}^{2,2}$ that does not come from $H_{\text{pr}}^{3,1}$

So the first module gives

$$c\left(\frac{h^3 + 3h^2 - 4h - 36}{6}\right) = c\left(\frac{b_2^3 - 3b_2^2 - 4b_2 - 24}{6}\right),$$

the second one $-d\left(\frac{h^2-h-4}{2}+2h+2\right)=d\left(\frac{h^2+3h}{2}\right)=d\left(\frac{b_2^2-b_2-2}{2}\right)$.

Definitely, in H^8 there are elements which come from those part of H^6 which is not generated by $\text{Sym}^3 H^2$ and two $\mathfrak{so}(b_2 + 2, \mathbb{C})$ -modules generated by elements of $H_{\text{pr}}^{3,1}$ and $H_{\text{pr}}^{2,2}$.

$$b_6 = \frac{(b_2 + 2)(b_2 + 1)b_2}{6} + c \left(\frac{b_2^2 - b_2 + 2}{2} \right) + db_2 + e$$

Then each element of $H^{3,3}$ part generate $\mathfrak{so}(b_2 + 2, \mathbb{C})$ -module of dimension $(b_2 + 2)$. That gives in b_8 the following term $e(b_2)$

$$b_8 \geq \frac{(b_2 + 3)(b_2 + 2)(b_2 + 1)b_2}{24} + c \left(\frac{b_2^3 - 3b_2^2 - 4b_2 - 24}{6} \right) + d \left(\frac{b_2^2 - b_2 - 2}{2} \right) + eb_2$$

From Salamon's relation we have

$$\begin{aligned} & 8 \cdot \frac{(b_2 + 2)(b_2 + 1)b_2}{6} + 8c \left(\frac{b_2^2 - b_2 + 2}{2} \right) + 8db_2 + 8e + 44 \left(\frac{(b_2 + 1)b_2}{2} + cb_2 + d \right) + \\ & \quad + 104b_2 + 188 + b_7 - 71b_3 - 23b_5 \geq \\ & \geq 2 \frac{(b_2 + 3)(b_2 + 2)(b_2 + 1)b_2}{24} + 2c \left(\frac{b_2^3 - 3b_2^2 - 4b_2 - 24}{6} \right) + 2d \left(\frac{b_2^2 - b_2 - 2}{2} \right) + 2eb_2 \end{aligned}$$

Then after rearranging we get

$$\begin{aligned} & -2 \frac{(b_2 + 3)(b_2 + 2)(b_2 + 1)b_2}{24} + 8 \cdot \frac{(b_2 + 2)(b_2 + 1)b_2}{6} + 44 \left(\frac{(b_2 + 1)b_2}{2} \right) + 104b_2 + 188 + b_7 \geq \\ & \geq c \left(\frac{b_2^3 - 15b_2^2 - 124b_2 - 48}{3} \right) + d(b_2^2 - 9b_2 - 46) + 2e(b_2 - 4) + 71b_3 + 23b_5 \end{aligned}$$

The left-part is $\frac{-b_2^4 + 10b_2^3 + 301b_2^2 + 530b_2 + 2256}{12} + b_7$.

Now recall that all odd Betti numbers are zero. Then the left-hand side is negative for $b_2 \geq 24$, although the right-hand side is positive. ■

Remark: The second Betti number b_2 is at most 24 for a sufficiently small b_7 . Indeed, for the proof of Theorem 4.1 we use the fact that

$$F(b_2) + b_7 := \frac{-2b_2^4 + 20b_2^3 + 602b_2^2 + 1060b_2 + 4512}{24} + b_7$$

is negative than $b_2 \geq 25$. This is also true for $b_7 \leq |F(b_2)|_{b_2=25} = 1281$.

Theorem 4.2: Let M be a ten-dimensional hyperkähler manifold with $H^{2k+1}(M, \mathbb{C}) = 0$. Then $b_2 \leq 25$.

Proof: The proof is very similar to the previous one. We could see that they are contributions from $H_{\text{pr}}^{3,1}, H_{\text{pr}}^{2,2}, H_{\text{pr}}^{3,3}, H_{\text{pr}}^{4,4}$. They are isomorphic to $\bigwedge^4 \mathbb{C}^{b_2+2}, \bigwedge^3 \mathbb{C}^{b_2+2}, \bigwedge^2 \mathbb{C}^{b_2+2}$, and \mathbb{C}^{b_2+2} as irreducible $\mathfrak{so}(b_2 + 2, \mathbb{C})$ -modules.

The explicit calculations like in the case of dimension eight give us the following

$$\frac{1}{60}(b_2 + 3)(b_2 + 4)(b_2 + 10)(-b_2^2 - 21b_2 + 118) = Q(b_2, c, d, e, f, g),$$

Polynomial $Q(b_2)$ is positive for $b_2 \geq 26$ since c, d, e, f, g are positive constants defined above (f sits in $H^{4,4}$ -part, g generates $H^{5,5}$). The left-hand side is negative for $b_2 \geq 26$. Hence, $b_2 \leq 25$. ■

References

- [B] F.A. Bogomolov, *On the decomposition of Kähler manifolds with trivial canonical class*, *Math. USSR-Sb.* **22**, pp. 580 - 583, 1974.
- [Bea] A. Beauville, Varieties Kahleriennes dont la première classe de Chern est nulle, *J. Diff. Geom.*, **16**, pp. 755-782, 1983.
- [Bea1] A. Beauville, Holomorphic symplectic geometry: a problem list. Preprint arXiv:1002.4321v1 [math.AG].
- [G] D. Guan, On the Betti numbers of irreducible compact hyperkähler manifolds of complex dimension four, *Math. Res. Lett.*, **8**, 5-6, pp 663-669, 2001.
- [H] D. Huybrechts, Finiteness results for hyperkähler manifolds, *J. Reine Angew. Math.*, **558**, pp. 1522, 2003. Preprint version arXiv:0109024 [math.AG].
- [HS] N. Hitchin and J. Sawon, Curvature and Characteristic Numbers of Hyperkähler Manifolds, *Duke Math. J.*, 2001, **106(3)**, 599-615.
- [K] N. Kurnosov, About inequality involving Betti numbers of hyperkahler a manifolds of dimension six, *Mathematical Notes*, **1**, 99, 2016.
- [LL] E. Looijenga and V. Lunts, A Lie algebra attached to a projective variety, *Invent. Math.*, **129**, no. 2, pp. 361-412, 1999.
- [O1] K.G. O’Grady, *Desingularized moduli spaces of sheaves on a K3*, *J. für die reine und angew. Math.*, **512**, pp. 49-117, 1999. Preprint arXiv:9708009v2 [math.AG].
- [O2] K.G. O’Grady, *A new six-dimensional irreducible symplectic variety*, *J. Algebraic Geom.*, **12**, pp. 435-505, 2003.
- [S] J. Sawon, *private communication*.
- [Sa] S. Salamon, On the cohomology of Kahler and hyper-Kahler manifolds, *Topology*, **35**, no. 1, pp. 137-155, 1996.
- [V1] M. Verbitsky, Actions of the Lie algebra of $SO(5)$ on the cohomology of a hyper-Kähler manifold, (English translation) *Funct. Anal. Appl.*, **24**, no. 3, pp. 229-230, 1990.

- [V2] M. Verbitsky, Cohomology of compact hyper-Kähler manifolds and its applications, *Geom. Funct. Anal.*, **6**, no. 4, pp. 601-611, 1995.
- [Y] S.T. Yau, *On the Ricci curvature of a compact Kähler manifold a and the complex Monge-Ampere equation I.*, *Comm. on Pure and Appl. Math.*, **31**, pp. 339-411, 1978.

NIKON KURNOSOV

LABORATORY OF ALGEBRAIC GEOMETRY AND ITS APPLICATIONS,
NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF ECONOMICS
7 VAVILOVA STR., MOSCOW, RUSSIA, 117312;
INDEPENDENT UNIVERSITY OF MOSCOW,
11 BOL.VLAS'EVSKIY PER., MOSCOW, RUSSIA, 119002
nikon.kurnosov@gmail.com